

# **Application Guidelines**

## **Storage & Handling:**

Do not freeze or store in direct sunlight. Do not store in temperatures above 80° F for extended periods of time.

## **Mixing & Application:**

Envita SC is compatible with most pesticides when in solution with water. Mixing order should be as follows:

### **In-furrow Applications:**

- 1) Add water to half of the application volume\*
- 2) Add tank mix partners\*\*
- 3) Shake bladder and add Envita SC as the last component
- 4) Agitate/circulate and apply as soon as possible
- 5) Apply with a minimum of 5 gpa total solution

### **Foliar Applications:**

- 1) Fill spray tank at least half-full of water\*
- 2) Add tank mix partners following label recommendations
- 3) Shake bladder and add Envita SC as the last component\*\*
- 4) Fill with water, agitate/circulate and apply as soon as possible
- 5) Apply with a minimum of 10 gpa total solution

Use rate: 0.8 fl. oz/acre
For use on multiple crops
In-furrow use: Apply directly to the seed, not 2x2.
Foliar application: Early Vegetative to Early Bloom - Refer to crop specific guidelines for primary application timings
Rain fast after 2 hours

Always follow tank mix partner application guidelines Ideally tank mix pH should be between 4 and 7 Optimal application temperatures:

- Above 80% relative humidity: 50 90°F
- Below 80% relative humidity: 50 80°F

OMRI certified

If you have further questions, please contact us at: info@envitasupport.com or call 1-877-ENVITA1 (1-877-368-4821).

Not all products are registered for sale in all states. Always read and follow label directions. Envita is a registered trademark of Azotic Technologies.









<sup>\*</sup>Envita SC is safe in chlorinated water up to 3.5 ppm

<sup>\*\*</sup>Use low salt-index fertilizers (SI <20) for in-furrow applications

<sup>\*</sup>Envita SC is safe in chlorinated water up to 3.5 ppm

<sup>\*\*</sup>If using Envita SC alone, use a non-ionic surfactant at a rate of 1 qt/100 gallons